



WEST BENGAL STATE UNIVERSITY

B.Sc. Honours 2nd Semester Examination, 2020

CEMACOR03T-CHEMISTRY (CC3)

Time Allotted: 2 Hours

Full Marks: 40

*The figures in the margin indicate full marks.
Candidates should answer in their own words and adhere to the word limit as practicable.
All symbols are of usual significance.*

Answer any four questions taking one from each unit

Unit-I

1. (a) Write down possible arrangements of electrons in p^3 configuration and identify the arrangement with maximum exchange energy. 2
- (b) What electronic transition in He^+ spectrum would have the same wavelength as the first Lyman transition of hydrogen? 3
- (c) How do the shapes of s and p orbitals can be obtained from angular function? Give reasons. 3
- (d) Show that frequency of revolution of an electron in Bohr orbit (quantum number n) is given by the expression. 3

$$\nu_{\text{orb}} = (4\pi^2 m z^2 e^4) / n^3 h^3$$

(Terms have their usual meaning)

Hence show that the frequency ν of the emitted radiation for transition from n_1 to n_2 , $(n_1 - n_2) = 1$ is intermediate between the frequencies of orbital revolution in these two orbits.

2. (a) Give the radial wave-function of the 3s hydrogenic orbital. How many radial nodes are there? 2+1
- (b) The velocity of an electron is 2×10^8 cm/sec. Calculate its wave length. 2
- (c) Stage the limitations of Aufbau principle with necessary illustrations. 3
- (d) Find out the spectroscopic ground state term symbols for Ti^{2+} and Co^{2+} ions. 3

Unit-II

3. (a) After calcium, electrons enter the 4s orbital before going to the 3d-orbitals, but when a transition metal ionizes, the 4s electrons are removed first. Why? 3
- (b) The electron affinity of Au is abnormally high and it may exist as auride — Justify. 2
- (c) Using Pauling's method, calculate the radii of K^+ and Cl^- ions. The observed $\text{K}^+ - \text{Cl}^-$ distance in KCl crystal is 314 pm. 3

4. (a) Calculate Z^* for the following electrons in a Scandium atom 3
 (i) 3p (ii) 3d (iii) 4s
 (b) Explain the variation of the second IE (kJ/mole) of the elements given in the parenthesis: Mg (1450), Al (1817), Si (1576), P (1903), S (2251), Cl (2297). 3
 (c) The atomic radii of Zr and Hf are almost identical — Explain. 2

Unit-III

5. (a) State solvent-system concept of acids and bases. Give one example of each of an acid and a base in liquid ammonia as solvent. 3
 (b) Why acidity in aqueous medium increases in the sequence 2

$$\text{CH}_4 < \text{NH}_3 < \text{H}_2\text{O} < \text{HF} ?$$

 (c) Why do Ca, Al and Ni exist in nature respectively as carbonate, oxide and sulphide? 3
 (d) SnCl_2 can act both as a Lewis acid and a Lewis base. Explain. 2
6. (a) What will be the order of acidity of H_3PO_4 , H_3PO_3 and H_3PO_2 ? Give reasons. 3
 (b) State the theory by which the reaction 2

$$6\text{CaO} + \text{P}_4\text{O}_{10} \rightarrow 2\text{Ca}_3(\text{PO}_4)_2$$
 may be regarded as acid-base reaction.
 (c) A buffer solution contains 0.10 mole of CH_3COOH and 0.10 mole of CH_3COO^- per litre. Calculate the pH of the buffer. [$K_a = 1.8 \times 10^{-5}$] 3
 (d) Arrange the given ions in order of increasing acidity in aqueous medium with justification. 2

$$[\text{Ni}(\text{H}_2\text{O})_6]^{2+}, [\text{Fe}(\text{H}_2\text{O})_6]^{3+}, [\text{Al}(\text{H}_2\text{O})_6]^{3+} \text{ and } [\text{Mn}(\text{H}_2\text{O})_6]^{2+}$$

Unit-IV

7. (a) Balance the following equation by Ion-electron Method 2

$$\text{Br}_2 + \text{NaOH} \rightarrow \text{NaBrO}_3 + \text{NaBr} + \text{H}_2\text{O}$$


 (b) According to reduction potential value of $\text{Cu}^{2+}/\text{Cu}^+$ ($E^0 = +0.15\text{V}$) and $\frac{1}{2} \text{I}_2/\text{I}^-$ ($E^0 = +0.54\text{V}$) system, Cu^{2+} should not oxidize I^- . Explain how can iodometric titration of Cu^+ be possible. [$K_{s(\text{CuI})} \approx 1 \times 10^{-2}$ at 25°C] 3
 (c) For a redox reaction 3

$$\text{MnO}_4^- + 5\text{Fe}^{2+} + 8\text{H}^+ \rightleftharpoons \text{Mn}^{2+} + 5\text{Fe}^{3+} + 4\text{H}_2\text{O}$$

 Calculate the equilibrium constant value
 (Given $E^0_{\text{MnO}_4^-/\text{Mn}^{2+}} = +1.52$ volt, $E^0_{\text{Fe}^{3+}/\text{Fe}^{2+}} = +0.77$ volt)
 (d) The solubility of AgCl is 0.0015 g dm^{-3} . Calculate its solubility product. 3
8. (a) What are the characteristics of redox indicators? Give one example of a redox indicator. 2+1
 (b) Give reason why Cl^- ion is oxidized to Cl_2 by KMnO_4 Solution at low pH. 2
 (Given $E^0_{\text{MnO}_4^-/\text{Mn}^{2+}} = +1.52$ volt, $E^0_{\frac{1}{2}\text{Cl}_2/\text{Cl}^-} = +1.36$ volt)

(c) What are disproportionation and comproportionation reactions? Give one example of each. 2+1

(d) From the following Latimer diagram 3



(i) Calculate the reduction potential of the reaction $\text{Sn}^{4+} \rightarrow \text{Sn}$.

(ii) Comment on the ease of reduction of Sn^{2+} to Sn and Sn^{4+} to Sn.

N.B. : *Students have to complete submission of their Answer Scripts through E-mail / Whatsapp to their own respective colleges on the same day / date of examination within 1 hour after end of exam. University / College authorities will not be held responsible for wrong submission (at in proper address). Students are strongly advised not to submit multiple copies of the same answer script.*

—x—